



Immediate Care Center

Date _____ Time _____ Account # _____ Rec _____

Name _____ DOB _____ Age _____ Sex _____ SSN _____

Street _____ City _____ State _____ Phone # _____

Do you have Medicare or Medicaid? Yes _____ No _____

- Influenza vaccine (A-H1N1+A-H3N2+B+B) is recommended for any person greater than 6 months of age who, because of age or underlying medical condition, is at risk for complications to influenza. Health care workers and others in contact with persons in the high risk groups should also be vaccinated. Influenza vaccine may also be administered to anyone who wishes to reduce the chance of becoming infected with the influenza virus. Influenza vaccine is propagated in eggs for preparation of the virus vaccine, therefore, a vaccine
➤ should not be administered to anyone with a history of allergies to eggs or egg products or thimerosal, anyone having an acute febrile illness or anyone with an acute neurological disorder.
➤ Because influenza vaccine contains only non-infectious viruses, it cannot cause influenza.
➤ Respiratory disease after vaccination represents coincidental illness unrelated to influenza vaccination. In children 6 months through 35 months of age, the most common (≥10%) injection-site reactions were pain (57%) or tenderness (54%), erythema (37%), and swelling (22%); the most common solicited systemic adverse reactions were irritability (54%), abnormal crying (41%), malaise (38%), drowsiness (38%), appetite loss (32%), myalgia (27%), vomiting (15%), and fever (14%). (6.1) • In children 3 years through 8 years of age, the most common (≥10%) injection-site reactions were pain (67%), erythema (34%), and swelling (25%); the most common solicited systemic adverse reactions were myalgia (39%), malaise (32%), and headache (23%). (6.1) • In adults 18 years and older, the most common (≥10%) injection-site reaction was pain (47%); the most common solicited systemic adverse reactions were myalgia (24%), headache (16%), and malaise (11%). (6.1) • In adults 65 years of age and older, the most common (≥10%) injection site reaction was pain (33%); the most common solicited systemic adverse reactions were myalgia (18%), headache (13%), and malaise.

Signature _____ Date _____

- Prevnar is indicated in children 6 weeks through 5 years of age (prior to the 6th birthday), Prevnar 13 is indicated for: • active immunization for the prevention of invasive disease caused by Streptococcus pneumoniae serotypes active immunization for the prevention of otitis media caused by S. pneumoniae serotypes No otitis media efficacy data are available for serotypes In children 6 years through 17 years of age (prior to the 18th birthday)
➤ Prevnar 13 is indicated for: • active immunization for the prevention of invasive disease caused by S. pneumoniae serotypes In adults 18 years of age and older,
➤ Prevnar 13 is indicated for: • active immunization for the prevention of pneumonia and invasive disease caused by S. pneumoniae serotypes Limitations of Prevnar 13 Use and Effectiveness •

In adults aged 18 years and older, the most commonly reported solicited adverse reactions (>5%) were pain at the injection site (>50%), fatigue (>30%), headache (>20%), muscle pain (>20%), joint pain (>10%), decreased appetite (>10%), injection site redness (>10%), injection site swelling (>10%), limitation of arm movement (>10%), vomiting (>5%), fever (>5%), chills (>5%), and rash (>5%).

Signature _____ Date _____

Mcare/Mcaid: Administer Flu Shot: G0008 \$15
Flu shot: 90688 \$10

Administer PN13 or PN23: G0009 \$25
PN13: 90670 \$200 PN23: 90732 \$90

Private Pay: Administer Flu Shot: 90471 \$15
Flu Shot (6-35 months): 90687 \$10
Flu Shot (3 AND UP): 90688 \$10

Administer PN13 or PN23: 90472 \$25
PN23: 90732 \$90 PN13: 90732 \$200

.5cc Flucelvax lot #195129 Exp 05/2018 in deltoid/thigh prepped with alcohol with 25g 1" needle without problems / ins:
.5cc PN lot# Exp in deltoid prepped with alcohol with 25g 1" needle without problems / ins:

Total: _____ Amount Paid: _____ How paid: Cash _____ CC _____ Check _____ Tax ID #: 82-0484708